

Note: Throughout this document, percents are rounded independently and may not add to 100%.

Summary Statistics of Reported and Verified Cases of Tuberculosis in San Joaquin County in 2006, (N=78)

I. Demographic Information

GENDER	NUMBER OF CASES	PERCENT OF CASES
Male	51	65.4%
Female	27	34.6%

AGE GROUP	NUMBER OF CASES	PERCENT OF CASES
0-4	3	3.8%
5-14	0	0.0%
15-24	6	7.7%
25-44	20	25.6%
45-64	24	30.8%
65+	25	32.1%

RACE	NUMBER OF CASES	PERCENT OF CASES
Hispanic	26	33.3%
Asian/P.I.	39	50.0%
Filipino	15	19.2%
Cambodian	7	9.0%
Indian	5	6.4%
Chinese	2	2.6%
Vietnamese	2	2.6%
Asian (unspecified)	2	2.6%
Loatian	1	1.3%
Hmong	1	1.3%
Japanese	1	1.3%
Pakistani	1	1.3%
Sri Lankan	1	1.3%
Burmese	1	1.3%
White	5	6.4%
Black	7	9.0%
Multi-race¹	1	1.3%

¹Multi-race individual is American Indian and White

Note: Throughout this document, percents are rounded independently and may not add to 100%.

**Summary Statistics of
Reported and Verified Cases of Tuberculosis in
San Joaquin County in 2006, (N=78)**

I. Demographic Information

CITY OF RESIDENCE & ZIP CODES	NUMBER OF CASES	PERCENT OF CASES
STOCKTON	58	74.4%
95202	3	3.8%
95203	14	17.9%
95204	2	2.6%
95205	10	12.8%
95206	11	14.1%
95207	2	2.6%
95209	3	3.8%
95210	9	11.5%
95212	2	2.6%
95215	1	1.3%
95219	1	1.3%
LODI	6	7.7%
95240	4	5.1%
95242	2	2.6%
TRACY	7	9.0%
95376	3	3.8%
95377	3	3.8%
95391	1	1.3%
MANTECA	4	5.1%
95336	2	2.6%
95337	2	2.6%
LATHROP (zip code = 95330)	2	2.6%
WOODBIDGE (zip code = 95258)	1	1.3%

Note: Throughout this document, percents are rounded independently and may not add to 100%.

Summary Statistics of Reported and Verified Cases of Tuberculosis in San Joaquin County in 2006, (N=78)

I. Demographic Information

COUNTRY OF ORIGIN	NUMBER OF CASES	PERCENT OF CASES
U.S. BORN	28	35.9%
FOREIGN BORN	50	64.1%
Philippines	16	20.5%
Mexico	11	14.1%
India	5	6.4%
Cambodia	5	6.4%
Vietnam	2	2.6%
China	2	2.6%
Laos	2	2.6%
Botswana	1	1.3%
Burma	1	1.3%
Pakistan	1	1.3%
Sri Lanka	1	1.3%
Cuba	1	1.3%
Peru	1	1.3%
Yemen	1	1.3%

II. Associated Risk Factors Among Reported Cases At The Time of Diagnosis

RISK FACTOR	NUMBER OF CASES	PERCENT OF CASES
Homelessness	16	20.5%
Imprisonment	0	0.0%
Long-term Care Institutionalization	3	3.8%
Injection Drug Use within Past Year	6	7.7%
Non-injection Drug Use within Past Year	10	12.8%
Excess Alcohol Use within Past Year	15	19.2%
Associated AIDS	4	5.1%

Note: Throughout this document, percents are rounded independently and may not add to 100%.

Summary Statistics of Reported and Verified Cases of Tuberculosis in San Joaquin County in 2006, (N=78)

III. Clinical Profile of Reported Cases

CLINICAL FACTOR	NUMBER OF CASES	PERCENT OF CASES
Previous Diagnosis of Tuberculosis	1	1.3%
Tuberculin Skin Test – Positive PPD	63	80.8%
Sputum Smear – Positive AFB	30	38.5%
Sputum Culture – Positive	47	60.3%
Chest X-ray - Normal	9	11.5%
Chest X-ray - Cavitory	16	20.5%
Chest X-ray - Noncavitory	51	65.4%
Major Site of Disease – Pulmonary	59	75.6%
Major Site of Disease – Extrapulmonary	15	19.2%
Pleural	4	5.1%
Lymphatic: Other	3	3.8%
Bone and/or joint	3	3.8%
Lymphatic: Cervical	2	2.6%
Peritoneal	1	1.3%
Genitourinary	1	1.3%
Other	1	1.3%
Major Site of Disease – Both Pulmonary and Extrapulmonary	4	5.1%

Note: The tuberculin skin test was not conducted in 2 (2.6%) cases. Both the sputum smear and the sputum culture tests were not done in 10 (12.8%) cases.

Note: Throughout this document, percents are rounded independently and may not add to 100%.

Summary Statistics of Reported and Verified Cases of Tuberculosis in San Joaquin County in 2006

IV. Anti-Tuberculosis Drug Sensitivity Profile (N=59)

MEDICATION	NUMBER OF CASES	PERCENT OF CASES
First Line Drugs – Tests Performed¹	59	100.0%
Sensitive to all first line drugs	53	89.8%
Resistant to at least one first line drug	6	10.2%
Isoniazid (INH) – Resistant	4	6.8%
Rifampin – Resistant	0	0.0%
Pyrazinamide (PZA) - Resistant	0	0.0%
Ethambutol – Resistant	0	0.0%
Streptomycin - Resistant	3	5.1%
Other Drugs² - Resistant	0	0.0%
Multi-Drug Resistant³	0	0.0%

¹There was one case in which only INH and Rifampin testing were done. There were also 6 cases in which Streptomycin was not tested.

²Other drugs tested on one case included Ethionamide, Kanamycin, and Capreomycin.

³Multidrug resistant definition is resistant to at least INH and Rifampin.